

MYCOTAXON

Volume 116, pp. 395–400

DOI: 10.5248/116.395

April–June 2011

Two new species of *Septobasidium* (Septobasidiaceae) from Gaoligong Mountains in ChinaCHUNXIA LU^{1,2} & LIN GUO^{1*}¹Key Laboratory of Systematic Mycology and Lichenology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China²Graduate University of Chinese Academy of Sciences, Beijing 100049, China

CORRESPONDENCE TO *: Ch.x.lu@hotmail.com & *guol@im.ac.cn

ABSTRACT —Two new species, *Septobasidium lyoniae* on *Lyonia ovalifolia* associated with *Pseudaulacaspis* sp. and *Septobasidium pittospori* on *Pittosporum kerrii* associated with *Chionaspis* sp., are described. They were collected from Gaoligong Mountains in Yunnan Province, China.

KEY WORDS —*Pucciniomycetes*, *Septobasidiales*, taxonomy

Previously, four new species of *Septobasidium* were found in Gaoligong Mountains of Yunnan Province (Lu & Guo 2009b, 2010a). From the same area an additional two new species are described as follows:

***Septobasidium lyoniae* C.X. Lu & L. Guo, sp. nov.**

FIGS. 1–7

MYCOBANK MB 519997

Basidiomata resupinata, 2–11 cm longa, 1–5 cm lata, cinnamomeo-brunnea, fumoso-brunnea vel brunnea, margine determinata, superficie laevia, in sectione 510–1200 µm crassa. Subiculum 25–50 µm crassum, brunneum, e subiculo stratum hypharum 275–580 µm altum vel columnam 200–250 altam, 110–130 µm latam formans. Ab strato hymenii hyphae saepe repullulantes tum stratum hypharum et hymenium secundum formantes. Stratum hymenii 100–220 µm altum, basidiis bistratos praeditum. Hyphae hymenii 3.5–5 µm crassae. Basidia cylindrica, recta vel leviter curvata, 4-cellularia, 45–57 × 8.5–10.5 µm, hyalina or brunneola. Sterigmata conica, 5–6 µm longa. Sine probasidio. Basidiosporae non visae. Haustoria ex hyphis irregulariter spiralibus constantia.

TYPE: On *Lyonia ovalifolia* (Wall.) Drude (*Ericaceae*): China, Yunnan Province, Gaoligong Mountains, Longling, alt. 1100 m, 6.IX.2008, S.H. He, Y.F. Zhu & L. Guo 2382, HMAS 250384 (**holotype**), associated with *Pseudaulacaspis* sp. (*Diaspididae*).

Basidiomata on branches, resupinate, perennial, 2–11 cm long, 1–5 cm wide, cinnamon-brown, smoke-brown or brown; margin determinate; surface smooth. In section 510–1200 µm thick. Subiculum 25–50 µm thick, brown.

From the subiculum forming the hyphal layer 275–580 μm thick, or pillars 200–250 μm high and 110–130 μm wide. From hymenial layer the fungal hyphae often renew growth to form the second hyphal layer and hymenium. The scale insects present between the hymenium and the hyphal layer. Hymenial layer 100–220 μm thick, usually with two continual basidial layers and upright hymenial hyphae 3.5–5 μm thick. Basidia arising directly from the hyphae without a probasidial cell, cylindrical, straight or slightly curved, 4-celled, 45–57 \times 8.5–10.5 μm , hyaline or brownish. Sterigmata coniform, 5–6 μm long. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.









FIG. 1. Basidia of *Septobasidium lyoniae* (HMAS 250384, holotype).

FIGS. 2–7. *Septobasidium lyoniae* (HMAS 250384, holotype). 2. Basidiomata on branch. 3–4. Sections of basidiomata (scale insects are arrowed). 5–6. Basidia (arrows). 7. Haustoria.



REMARKS: Morphologically, *Septobasidium lyoniae* is similar to *S. cirratum* Burt, from which it differs in producing 2-stratose hymenia (single layer in *S. cirratum*), and having thicker and non-coiled hymenial hyphae (3.5–5 μm vs 1.5–2 μm).




FIG. 8. Basidia of *Septobasidium pittospori* (HMAS 240137, holotype).

***Septobasidium pittospori* C.X. Lu & L. Guo, sp. nov.**

FIGS. 8–14






MYCOBANK MB 519998

Basidiomata resupinata, 13–20 cm longa, 2–5 cm lata, cinnamomeo-brunnea, margine determinata, superficie laevia, in vetustate caduca, in sectione 490–920 μm crassa. Subiculum brunneum, 30–55 μm crassum. Columnae brunneae, 190–290 μm altae, 25–190 μm crassae. Stratum hypharum 340–580 μm altum. Hymenium 50–90 μm crassum. Basidia cylindrica, recta vel leviter curvata, 4-cellularia, 35–50 \times 7–11 μm , hyalina or flavido-brunnea. Sine probasidio. Basidiosporae non visae. Haustoria ex hyphis et cellulis globularibus constantia.

TYPE: On *Pittosporum kerrii* Craib (*Pittosporaceae*): China, Yunnan Province, Gaoligong Mountains, Baoshan, Baihualin, alt. 1400 m, 3.IX.2008, S.H. He, Y.F. Zhu & L. Guo 2305, HMAS 240137 (holotype), associated with *Chionaspis* sp. (*Diaspididae*).

Basidiomata on branches, resupinate, 13–20 cm long, 2–5 cm wide, cinnamon-brown; margin determinate; surface smooth, peeling off partly in old stage. In section 490–920 μm thick. Subiculum brown, 30–55 μm thick. Pillars brown, 190–290 μm high, 25–190 μm wide. Hyphal layer 340–580 μm high. Hymenial layer 50–90 μm thick, having loosely and irregularly upright hymenial hyphae.

FIGS. 9–14. *Septobasidium pittospori* (HMAS 240137, holotype). 9. Basidiomata on branches. 10–11. Sections of basidiomata. 12–13. Basidia (arrows). 14. Haustoria.



Basidia arising directly from the hyphae without a probasidial cell, cylindrical, straight or slightly curved, 4-celled, $35\text{--}50 \times 7\text{--}11 \mu\text{m}$, hyaline or yellowish brown. Basidiospores not seen. Haustoria consisting of hyaline hyphae and globose cells.

REMARKS: Morphologically, *Septobasidium pittospori* is similar to *S. lichenicola* (Berk. & Broome) Petch, from which it differs in having distinct pillars; *S. lichenicola* has inconspicuous pillars.

To date, 31 species of *Septobasidium* have been reported in China (Sawada 1933, Couch 1938, Teng 1963, Tai 1979, Kirschner & Chen 2007, Lu & Guo 2009a,b,c, 2010a,b,c, Lu et al. 2010, Chen & Guo 2011), including the two new species reported in this paper.

Acknowledgements

The authors would like to express their deep thanks to Dr. Eric H.C. McKenzie (Auckland, New Zealand) and Shuanghui He (Beijing Forestry University) for serving as pre-submission reviewers, to Dr. Shaun Pennycook (Auckland, New Zealand) for nomenclatural review, to Prof. Jianyun Zhuang (Institute of Microbiology, Chinese Academy of Sciences) for Latin corrections, to Prof. Zhenyu Li (Institute of Botany, Chinese Academy of Sciences) for identifying the host plants, to Prof. Sanan Wu (Beijing Forestry University) for identifying the scale insects, and to Mrs. Xiangfei Zhu for inking in line drawings. This study was supported by the National Natural Science Foundation of China (No. 30499340 and No. 30670005).

Literature cited

- Chen SZ, Guo L. 2011. *Septobasidium sichuanense* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 115: 481–484. doi: 10.5248/115.481
- Couch JN. 1938. The genus *Septobasidium*. Univ. of North Carolina Press, Chapel Hill. 480 p.
- Kirschner R, Chen CJ. 2007. New reports of two hypophyllous *Septobasidium* species from Taiwan. Fung. Sci. 22(1,2): 39–46.
- Lu CX, Guo L. 2009a. *Septobasidium maesae* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 109: 103–106. doi:10.5248/109.103
- Lu CX, Guo L. 2009b. Two new species of *Septobasidium* (*Septobasidiaceae*) from China. Mycotaxon 109: 477–482. doi:10.5248/109.477
- Lu CX, Guo L. 2009c. *Septobasidium annulatum* sp. nov. (*Septobasidiaceae*) and *Septobasidium kameii* new to China. Mycotaxon 110: 239–245. doi:10.5248/110.239
- Lu CX, Guo L. 2010a. Three new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. Mycotaxon 112: 143–151. doi:10.5248/112.143
- Lu CX, Guo L. 2010b. Two new species of *Septobasidium* (*Septobasidiaceae*) and *S. pallidum* new to China. Mycotaxon 113: 87–93. doi: 10.5248/113.87
- Lu CX, Guo L. 2010c. Two new species of *Septobasidium* (*Septobasidiaceae*) from Hainan province in China. Mycotaxon 114: 217–223. doi: 10.5248/114.217
- Lu CX, Guo L, Wei JG, Li JB. 2010. Two new species of *Septobasidium* (*Septobasidiaceae*) from southern China. Mycotaxon 111: 269–274. doi:10.5248/111.269
- Sawada K. 1933. Descriptive catalogue of the Formosan fungi. Part VI. Rep. Dept. Agric. Govt. Res. Inst. Formosa 61: 1–99.
- Tai FL. 1979. Sylloge Fungorum Sinicorum. Science Press, Beijing. 1527 p.
- Teng SC. 1963. Fungi of China. Science Press, Beijing. 808 p.